

CHAPTER I
CONSERVATION OF CONTINUOUS
QUANTITIES

EVERY notion, whether it be scientific or merely a matter of common sense, presupposes a set of principles of conservation, either explicit or implicit. It is a matter of common knowledge that in the field of the empirical sciences the introduction of the principle of inertia (conservation of rectilinear and uniform motion) made possible the development of modern physics, and that the principle of conservation of matter made modern chemistry possible. It is unnecessary to stress the importance in every-day life of the principle of identity; any attempt by thought to build up a system of notions requires a certain permanence in their definitions. In the field of perception, the schema of the permanent object¹ presupposes the elaboration of what is no doubt the most primitive of all these principles of conservation. Obviously conservation, which is a necessary condition of all experience and all reasoning, by no means exhausts the representation of reality or the dynamism of the intellectual processes, but that is another matter. Our contention is merely that conservation is a necessary condition for all rational activity, and we are not concerned with whether it is sufficient to account for this activity or to explain the nature of reality.

This being so, arithmetical thought is no exception to the rule. A set or collection is only conceivable if it remains unchanged irrespective of the changes occurring in the relationship between the elements. For instance, the permutations of the elements in a given set do not change its value. A number is only intelligible if it remains identical with itself, whatever the distribution of the units of which it is composed. A continuous quantity such as a length or a volume can only be used in reasoning if it is a permanent whole, irrespective of the possible arrangements of its parts. In a word, whether it be a matter of continuous or discontinuous qualities, of quantitative relations perceived in the sensible universe, or of sets and numbers conceived by thought, whether it be a matter of the child's earliest contacts with number or of the most refined axiomatizations of any intuitive system, in

¹ *La Construction du Réel chez l'Enfant*, chapter i.

each and every case the conservation of something is postulated as a necessary condition for any mathematical understanding.

From the psychological point of view, the need for conservation appears then to be a kind of functional *a priori* of thought. But does this mean that arithmetical notions acquire their structure because of this conservation, or are we to conclude that conservation precedes any numerical or quantifying activities, and is not only a function, but also an *a priori* structure, a kind of innate idea present from the first awareness of the intellect and the first contact with experience? It is experiment that will provide the answer, and we shall try to show that the first alternative is the only one that is in agreement with the facts.

§1. Technique and general results

This chapter and the one that follows will be devoted to experiments made simultaneously with continuous and discontinuous quantities. It seemed to us essential to deal with the two questions at the same time, although the former are not arithmetical and were to be treated separately in a special volume,¹ since it was desirable to ascertain that the results obtained in the case of discontinuous sets were general.

The child is first given two cylindrical containers of equal dimensions (A_1 and A_2) containing the same quantity of liquid (as is shown by the levels). The contents of A_2 are then poured into two smaller containers of equal dimensions (B_1 and B_2) and the child is asked whether the quantity of liquid poured from A_2 into ($B_1 + B_2$) is still equal to that in A_1 . If necessary, the liquid in B_1 can then be poured into two smaller, equal containers (C_1 and C_2), and in case of need, the liquid in B_2 can be poured into two other containers C_3 and C_4 identical with C_1 and C_2 . Questions as to the equality between ($C_1 + C_2$) and B_2 , or between ($C_1 + C_2 + C_3 + C_4$) and A_1 , etc., are then put. In this way, the liquids are subdivided in a variety of ways, and each time the problem of conservation is put in the form of a question as to equality or non-equality with one of the original containers. Conversely, as a check on his answers, the child can be asked to pour into a glass of a different shape a quantity of liquid approximately the same as that in a given glass, but the main problem is still that of conservation as such.

The results obtained seem to prove that continuous quantities are not at once considered to be constant, and that the notion of conservation is gradually constructed by means of an intellectual

¹ J. Piaget and B. Inhelder, *Le Développement des Quantités chez l'Enfant (Conservation et Atomisme)*, 1941.

mechanism which it is our purpose to explain. By grouping the answers to the various questions, it is possible to distinguish three stages. In the first, the child considers it natural for the quantity of liquid to vary according to the form and dimensions of the containers into which it is poured. Perception of the apparent changes is therefore not corrected by a system of relations that ensures invariance of quantity. In the second stage, which is a period of transition, conservation gradually emerges, but although it is recognized in some cases, of which we shall attempt to discover the characteristics, it is not so in all. When he reaches the third stage, the child at once postulates conservation of the quantities in each of the transformations to which they are subjected. Naturally this does not mean that this generalization of constancy extends at this stage beyond the limits of the field studied here.

In our interpretation of these facts, we can start from the following hypotheses, some of which directed the research of this chapter while others arose in the course of our experiments. The question to be considered is whether the development of the notion of conservation of quantity is not one and the same as the development of the notion of quantity. The child does not first acquire the notion of quantity and then attribute constancy to it; he discovers true quantification only when he is capable of constructing wholes that are preserved. At the level of the first stage, quantity is therefore no more than the asymmetrical relations between qualities, i.e., comparisons of the type 'more' or 'less' contained in judgements such as 'it's higher', 'not so wide', etc. These relations depend on perception, and are not as yet relations in the true sense, since they cannot be co-ordinated one with another in additive or multiplicative operations. This co-ordination begins at the second stage and results in the notion of 'intensive' quantity, i.e., without units, but susceptible of logical coherence. As soon as this intensive quantification exists, the child can grasp, before any other measurement, the proportionality of differences, and therefore the notion of extensive quantity. This discovery, which alone makes possible the development of number, thus results from the child's progress in logic during these stages.

§2. Stage I: Absence of conservation

For children at the first stage, the quantity of liquid increases or diminishes according to the size or number of the containers. The reasons given for this non-conservation vary from child to child, and from one moment to the next, but (in every case the child thinks that the change he sees involves a change in the total value of the liquid.) Here we have some examples:

CONSERVATION OF QUANTITIES

Blas (4;0). 'Have you got a friend?—*Yes, Odette.*—Well look, we're giving you, Clairette, a glass of orangeade (A1, $\frac{3}{4}$ full), and we're giving Odette a glass of lemonade (A2, also $\frac{3}{4}$ full). Has one of you more to drink than the other?—*The same.*—This is what Clairette does: she pours her drink into two other glasses (B1 and B2, which are thus half filled). Has Clairette the same amount as Odette?—*Odette has more.*—Why?—*Because we've put less in.* (She pointed to the levels in B1 and B2, without taking into account the fact that there were two glasses).—(Odette's drink was then poured into B3 and B4.) *It's the same.*—And now (pouring Clairette's drink from B1 + B2 into L, a long thin tube, which is then almost full)?—*I've got more.*—Why?—*We've poured it into that glass* (pointing to the level in L), *and here* (B3 and B4) *we haven't.*—But were they the same before?—*Yes.*—And now?—*I've got more.* Clairette's orangeade was then poured back from L into B1 and B2: 'Look, Clairette has poured hers like Odette. So, is all the lemonade (B3 + B4) and all the orangeade (B1 + B2) the same?—*It's the same* (with conviction).—Now Clairette does this (pouring B1 into C1 which is then full, while B2 remains half full). Have you both the same amount to drink?—*I've got more.*—But where does the extra come from?—*From in there* (B1).—What must we do so that Odette has the same?—*We must take that little glass* (pouring part of B3 into C2).—And is it the same now, or has one of you got more?—*Odette has more.*—Why?—*Because we've poured it into that little glass* (C2).—But is there the same amount to drink, or has one got more than the other?—*Odette has more to drink.*—Why?—*Because she has three glasses* (B3 almost empty, B4 and C2, while Clairette has C1 full and B2).'

A moment later, a new experiment. Clairette was again shown glasses A1 and A2, $\frac{3}{4}$ full, one with orangeade for herself and the other with lemonade for Odette. 'Are they exactly the same?—*Yes* (verifying the levels).—Well, Odette is going to pour hers (A2) into all those (C1, C2, C3, C4, which were thus about half full). Have you both the same amount?—*I've got more. She has less. In the glasses there's less* (looking carefully at the levels).—But before, you both had the same?—*Yes.*—And now?—*Here* (pointing to the level in A1) *it's more, and here* (indicating the 4 glasses C) *it's less.*

Finally she was given only the big glass A1 almost full of orangeade: 'Look, Clairette does this: she pours it like that (into B1 and B2, which are then $\frac{3}{4}$ full). Is there more to drink now than before, or less, or the same?—*There's less* (very definitely).—Explain to me why.—*When you poured it out, it made less.*—But don't the little glasses together make the same?—*It makes less.*

Sim (5;0). She was shown A1 and A2 half full. 'There's the same amount in the glasses, isn't there?—(She verified it) *Yes.*—Look, Renée, who has the lemonade, pours it out like this (pouring A1 into B1 and B2, which were thus about $\frac{3}{4}$ full). Have you both still the same amount to drink?—*No. Renée has more because she has two glasses.*—What could you do to have the same amount?—*Pour mine into two glasses.* (She poured A2 into B3 and B4.)—Have you both got the same now?—(She looked for a long time at the 4 glasses *Yes.*—Now Madeleine (herself)

CONSERVATION OF CONTINUOUS QUANTITIES 7

is going to pour her two glasses into three (B3 and B4 into C1, C2 and C3). Are they the same now?—*No.*—Who has more to drink?—*Madeleine, because she has three glasses. Renée must pour hers too into three glasses.* (Renée's B1 and B2 were poured into C5, C6 and C7). There. —*It's the same.*—But now Madeleine pours hers into a fourth glass (C4, which was filled with a little from C1, C2 and C3). Have you both the same amount?—*I've got more.*—Is there more of the lemonade (C5, C6 and C7) or of the orangeade (C1, C2, C3 and C4)?—*The orangeade.*—(The two big glasses A1 and A2 were then put before her.) Look, we're going to pour back all the lemonade into this one (A1) as it was before, and all the orangeade into that one. Where will the lemonade come up to?—(She indicated a certain level)—And the orangeade?—(She indicated a higher level.)—Will the orangeade be higher than the lemonade?—*Yes, there's more orangeade* (pointing to the level she had indicated) *because there's more orangeade here* (pointing to C1, C2, C3 and C4).—You think it will come up to here?—*Yes.*—(This level was marked by an elastic band and she herself poured in the liquid and was delighted to find that it came up to the band. But when she poured the lemonade into A1 she was very much surprised to find that it reached the same level.) *It's the same!*—How's that?—*I think we've put a little back, and now it's the same.*

It is clear that so far the child had thought that there were changes in quantity when the number of glasses varied, but with the next question the level intervenes: 'Look, Madeleine pours the orangeade into that glass (A2 was poured into L, which was longer and narrower. L was then $\frac{3}{4}$ full, whereas the lemonade in A1 came only half way up.)—*There's more orangeade, because it's higher.*—Is there more to drink, or does it just look as if there is?—*There's more to drink.*—And now (pouring the lemonade into B1 and B2 and the orangeade into D1 and D2 which were wide, low glasses)?—*It's the orangeade that's more, because there* (in D1 and D2) *there's a lot.*—So if we pour the lemonade and the orangeade back here (A2 and A1), will the orangeade come up higher or will they be the same?—*Higher.*' She poured D1 and D2 back into A2, and B1 and B2 back into A1, and was again much surprised to see that the levels were the same.

Lac (5;6). 'Here are two glasses (A1 half full of orangeade and A2 slightly less full of lemonade.) The orangeade is for you and the lemonade for Lucien. Lucien is cross because he has less. He pours his drink into these two glasses (pouring A2 into B1 and B2). Who has more?—(Lac looked at the levels) *Me.*—Now you pour your drink into these two glasses (B3 and B4, the levels being thus slightly higher than in B1 and B2). Who has more?—*Me.*—And now Lucien takes this glass (B1) and divides it between these two (C1 and C2, which are then full, whereas B2 remains half-full). Who has more?—(Lac compared the levels and pointed to glasses C) *Lucien.*—Why?—*Because the glasses get smaller* (and therefore the levels rise).—But how did that happen? Before it was you who had more and now it's Lucien?—*Because there's a lot.*—But how did it happen?—*We took some.*—But where?—...—And how?—...—Has one of you got more?—*Yes, Lucien* (very definitely).

—And if I pour all the orangeade and all the lemonade into the two big glasses (A₁ and A₂) who will have more?—*I shall* (thus showing that he remembered the original position).—Then where has the extra you had gone to?—...—What could you do to have the same amount as Lucien? You can use any of the glasses.—Lac then took B₃ and poured some of it into C₃, an empty glass. He filled it, and put it opposite Lucien's C₁ and C₂. Then he compared B₃ to Lucien's B₂ and saw that there was less in B₃ than in B₂. He then took C₃ again, poured it back into B₃, and then, showing great disappointment, cried: *But why was it quite full there (C₃), and now (B₃) it isn't full any longer?*

Mus (5;0). This child, like those quoted earlier, relied on the number of glasses or the level, but in her case as in several others there was also a new factor, the size of the glasses. Nevertheless she followed three successive lines of thought:

I. *Size of the containers.*—She was given A₁ and A₂, $\frac{3}{4}$ full: 'Is there the same amount in both of them?'—*Yes.*—Olga pours hers out like this (A₂ into B₁ and B₂, almost full). Has she still the same amount?—*No.*—Who has more to drink?—*Gertrude (A₁).*—Why?—*Because she has a bigger glass.*—How is it that Olga has less?—...—And if I pour these (B₁ and B₂) back into that one (A₂) how will it be?—*The same amount (as in A₁).*—(I did so.) And if Olga pours it back again like this (A₂ into B₁ and B₂, almost full) is it the same?—*No.*—Why?—*It makes less.*

II. *Level.*—'Now Gertrude pours hers like this (A₁ into C₁ and C₂, almost filling C₁ and C₂ and leaving A₁ $\frac{1}{2}$ full). Who has more, Gertrude with those (A₁ + C₁ + C₂), or Olga with those (B₁ and B₂)?'—(She looked at the levels, which were about equal) *Both the same.*—Olga pours some of hers into another glass (B₃, thus lowering the general level in her glasses).—*Gertrude will have more. Olga will have less.*—Olga pours again into these glasses (B₁ and B₂ into C₃ and C₄, which were then full).—*She will have more (level).*—But before she had less; has she more now?—*Yes.*—Why?—*Because we put back here (C₃ and C₄) what was in the big glasses (B₁ and B₂).' The reasoning here was thus just the opposite of what it was in I.*

III. *Number of glasses and level.*—'If I give you some coffee in one cup, will it still be the same if you pour it into two glasses?'—*I'll have a little more.*—Where?—*In the two glasses of course.*—Mummy gives you two glasses of coffee (B₁ and B₂). Then you pour that one (B₂) into those (C₁ and C₂).—*There's more there (C₁ and C₂): there are two glasses quite full. There, there's only one.*—And of those (B₁ and the 4 C) which would you rather have, that one (B₁) or all those (4 C)?—*The big one (B₁).*—Why?—*Because there's more: the glass is big.*

Such then are the earliest reactions of the child confronted with the problem of conservation of quantities. He is not prepared to believe that a given quantity of liquid remains the same irrespective of changes in shape when it is poured from one container to another.

It might of course be argued that the child may not really have

grasped the question. Does he always understand that it refers to the total quantity, or does he think he is merely being asked about changes in the number, level or size of the glasses? But the problem is precisely to discover whether the child is capable of grasping a quantity as being a whole, as a result of the co-ordination of the various relationships he perceives. The fact that these children isolated one of these relationships may therefore be due as much to lack of understanding of the notions in question as to failure to grasp the verbal question.

On the other hand, it might be suggested that when the liquid is poured from one container to another before the eyes of the child there are certain illusions of perception that counteract his judgement as to conservation.¹ We are well aware that perception of the quantifiable qualities such as length, weight, etc., leads to systematic distortions, and that the child finds it extremely difficult to perceive the constancy of these qualities. Hence, when the constancy is directly perceived, there is no problem as far as we are concerned. Our only problem is to discover by what means the mind succeeds in constructing the notion of constant quantity in spite of the indications to the contrary provided by immediate perception. Judgement comes into play precisely when perception proves inadequate, and only then. For instance, the discovery that a given quantity of liquid does not vary when poured from a container A into one or two containers B of a different shape, requires on the part of the child an effort of intellectual understanding which will be the greater and the more easily analysable the more deceptive the immediate perception. We are therefore not concerned to discover why this perception is deceptive, but why children at a certain level accept it without question, whereas others correct it by the use of intelligence. Moreover, either perception must be studied 'from the angle of the object', in which case intelligence will in the final resort be the origin of the constancy, or else perception presupposes an organization which elaborates the constancy on its own plane, in which case the functioning and the successive structures of perception imply a sensory-motor activity that is intelligent from the start. If the latter is the case, the development of the notion of invariant quantities (like that of 'object') would be a continuation, on a new, abstract plane, of the work already undertaken by sensory-motor intelligence in the field of conservation of the object.

We shall attempt to interpret the examples given above from this second point of view. What is most striking at this first stage is the inadequate quantification of the perceived qualities, and the lack of co-ordination between the quantitative relations involved

¹ E. Brunsvik, *Wahrnehmung und Gegenstandswelt*. Leipzig u. Wien, 1939.

in the perception. For example, Blas (4;0) begins by thinking that the quantity of liquid diminishes when the contents of a large glass three-quarters full are poured into two smaller glasses, but that it increases when poured from these small glasses into a long, narrow tube. It is therefore only the level and not the number or the cross section of the glasses which seems to be Blas's criterion. But a moment later he thinks there is more liquid in three small glasses than in two medium-sized ones filled with the same quantity. There are two noteworthy features in this reaction. In the first place, the child continually contradicts himself. At one moment he thinks there is more orangeade than lemonade, at another he thinks the opposite, and yet it does not occur to him to question his previous assumption. Obviously, if it is accepted that a liquid is capable of expansion or contraction and has no constancy, there is no contradiction. The real contradiction lies in the fact that the child attempts to justify his opposing statements by resorting to explanations that he cannot co-ordinate one with another, and that lead to incompatible statements. Thus Blas sometimes finds his evidence in the level of the liquid and thinks that the quantity diminishes when it is poured from a large glass into several small ones: sometimes he bases his statement on the number of glasses involved, in which case the same operation is thought to imply an increase in quantity. Alternatively, the child will use the cross section of the containers in his estimate of the change, disregarding the number of glasses and the level, and will then take one of these factors into account and arrive at the opposite conclusion. This brings us to the second feature of the reaction: the child behaves as though he had no notion of a multi-dimensional quantity and could only reason with respect to one dimension at a time without co-ordinating it with the others. What has been said is true not only of Blas, but of all the children quoted above.

The reactions of this stage can therefore be interpreted in the following way. We must first look for the principle of differentiation between quantity and quality, and this from the first perceptual contact with the object. In every case, perception and concrete judgement attribute qualities to objects, but they cannot grasp these qualities without thereby relating them one to another. These relations can only be of two kinds: symmetrical relations expressing resemblances, and asymmetrical relations expressing differences. Now resemblances between qualities can only result in their classification (e.g., glasses C₁, C₂, C₃ . . . are 'equally small'), whereas asymmetrical differences imply 'more' and 'less' and thus indicate the beginnings of quantification (e.g., 'A₁ is larger than B₁' or 'A₁ is narrower than D'). In its primitive form, therefore, quantity is given at the same time as quality,

since it is constituted by the asymmetrical relations that necessarily link any qualities one to another. Qualities *per se* do not in fact exist; they are always compared and differentiated, and this differentiation, since it includes asymmetrical relations of differences, is the germ of quantity. From this point of view, the judgements characteristic of the first stage are obviously already quantitative in this sense. For instance, when Sim says: 'There's more orangeade because it's higher', she is merely expressing in terms of quantity a perceptual relation of difference between two qualities (the heights of the liquids).

At this first level, however, which we shall call that of 'gross quantity', quantification is restricted to the immediate perceptual relationships, just as 'gross quality' (i.e., directly perceived quality) is incapable of giving rise to a complete classification. The relationship of similarity between qualities will of course eventually result in a system of classes, but this only becomes possible with the elaboration of sequences of hierarchical inclusions involving the whole logic of classes and asymmetrical relations. As for the relations of difference, with which alone we are for the moment concerned, they will lead to a systematic quantification whose stages we shall study in subsequent chapters. But before this is achieved they must be able to satisfy two conditions that are lacking at this level, which accounts for the absence of measurable quantity and conservation.

The first of these conditions is that, from being mere perceptual relationships, they shall become true relations, thus giving rise to systems of graduations or 'intensive quantities'. (See Glossary.) Obviously a perceptual relationship does not as such constitute a relation. The criterion for the psychological existence of relations is the possibility of their composition, or in other words, the construction of their logical transitivity (or, if they cannot become transitive, the justification for their non-transitivity). The main characteristic of the perceptual relationships of gross quantity used by the child at this first level is that they cannot be composed one with another either additively or multiplicatively. When the child thinks that the quantity increases because the level rises, he is disregarding the cross section, and when he takes the cross section into account he disregards the level, and so on.¹

The following experiment makes this plain. The child is given

¹ By addition of asymmetrical relations we mean their actual or virtual seriation and the resulting graduation of the seriated terms. By multiplication of these relations we mean their seriation from two or more points of view simultaneously. In the examples quoted above, the simple series do not appear, but the children had to compare two quantities from several points of view, height, cross-section, number of glasses, etc., which constitutes multiplication of relations.

two containers A and L of equal height, A being wide and L narrow. A is filled to a certain height (one-quarter or one-fifth) and the child is asked to pour the same quantity of liquid into L. The dimensions are such that the level will be four times as high in L as in A for the same amount of liquid. In spite of this striking difference in the proportions, the child at this stage proves incapable of grasping that the smaller diameter of L will require a higher level of liquid. Those children who are clearly still at this stage are satisfied that there is 'the same amount to drink' in L when they have filled it to the same level as A.

Blas (4;0): 'Look, your mummy has poured out a glass of lemonade for herself (A) and she gives you this glass (L). We want you to pour into your glass as much lemonade as your mummy has in hers.—(She poured rather quickly and exceeded the level equal to that in A that she was trying to achieve.)—Will you both have the same like that?—*No.*—Who will have more?—*Me.*—Show me where you must pour to so that you both have the same.—(She poured up to the same level.)—Will you and mummy have the same amount to drink like that?—*Yes.*—Are you sure?—*Yes.*—Now watch what I'm doing (putting L' next to L). I'm going to pour that one (A) into this one (L'). Will that make the same here (L') as there (L)?—*Yes.*—(When I did so, the child laughed): *Mummy has more.*—Why?—...

Mus (5;0): 'Look (same story as for Blas). Show me with your finger how far I must pour.—*There* (indicating the same level in L as in A).—(I filled it slightly higher). Will there be the same amount to drink?—*You've put too much. There's a little more there* (in L). *I've a little more to drink*—What could you do to see if it's the same? (putting L' next to L).—...—Where will it come up to if we pour that one (A) into this one (L')?—*To there* (pointing to the same level as in A).—(I did so).—*Mummy has more* (with great surprise).—How did that happen?—*Because the glass (L') is smaller.* (Mus thus appeared to have grasped the relation height \times cross-section, but it was only a momentary glimpse, as we shall see.)—And if I pour this (L') back into that (A), which will have the most?—*Both a little, both the same.*—(I poured it back). Who has more to drink?—*Both less.*'

These reactions show that the child at the first stage is unable to reckon simultaneously with the height and cross section of the liquids he has to compare. It is not that he fails to notice the width of glass A when circumstances oblige him to make the comparison (e.g. Mus, when A is poured into L'), but when he merely has to estimate the quantities in A and L' he takes into account only the height.

The child at this stage has therefore not yet acquired the notion of multi-dimensional quantity, owing to lack of composition between the relationships of differences. For him the quantity of liquid does not depend on the combination of the various relations

of height, cross section, number of glasses, etc., since each of these relations is considered separately, as though independent of the others. Each relation therefore constitutes merely a 'gross quantity' that is essentially uni-dimensional. Even when the child uses terms such as 'big' or 'large' this quality is still, as the case of Mus shows, merely perceptual data not susceptible of composition with others, and therefore again constituting a uni-dimensional 'gross quantity'.

The second condition to be satisfied before these perceptual relations can lead to true quantification, namely that there shall be partition into equal units, is even more impossible of fulfilment at this first stage than is intensive graduation. Before there can be acceptance of the notion of conservation of the liquid, and therefore construction of the notion of extensive quantity (see Glossary), there must be understanding that every increase in height is compensated by a diminution in width, these two qualities being inversely proportional. Yet even in the very simple problem of the increase in the number of glasses, children at this stage show clearly that they are unable to grasp the fact that a quantity of liquid poured from one glass into two or three smaller glasses remains the same. Composition by partition is therefore as impossible as by relations.

§3. Stage II. Intermediary reactions

Between the children who fail to grasp the notion of conservation of quantity and those who assume it as a physical and logical necessity, we find a group showing an intermediary behaviour (not necessarily found in all children) which will characterize our second stage. Two at least of these transitional reactions are worthy of note. The first of these shows that the child is capable of assuming that the quantity of liquid will not change when it is poured from glass A into two glasses B₁ and B₂, but when three or more glasses are used he falls back on to his earlier belief in non-conservation. The second reaction is that of the child who accepts the notion of conservation when the differences in level, cross section, etc., are slight, but is doubtful when they are greater. Here we have some examples of the first type:

Edi (6;4): 'Is there the same in these two glasses (A₁ and A₂)?—*Yes.*—Your mummy says to you: Instead of giving you your milk in this glass (A₁), I give it to you in these two (B₁ and B₂), one in the morning and one at night. (It is poured out.) Where will you have most to drink, here (A₂) or there (B₁ + B₂)?—*It's the same.*—That's right. Now, instead of giving it to you in these two (B₁ and B₂), she gives it to you in three (pouring A₂ into C₁, C₂ and C₃), one in the morning, one at

lunch-time and one at night. Is it the same in the two as in the three, or not?—*It's the same in 3 as in 2 . . . No, in 3 there's more.*—Why? . . . —(B₁ and B₂ were poured back into A₁.) And if you pour the three (C₁ + C₂ + C₃) back into that one (A₂) how far up will it come?—(He pointed to a level higher than that in A₁.)—And if we pour these 3 into 4 glasses (doing so into C₁ + C₂ + C₃ + C₄, with a consequent lowering of the level) and then pour it all back into the big one (A₂), how far up will it come?—(He pointed to a still higher level.)—And with 5?—(He showed a still higher level.)—And with 6?—*There wouldn't be enough room in the glass.*

Pie (5;0): 'Is there the same amount here (A₁) and there (A₂)?—(He tested the levels.) *Yes.*—(A₁ was poured into B₁ + B₂). Is there the same amount to drink in these two together as in the other?—(He examined the levels in B₁ and B₂, which were higher than in A₁.) *There's more here.*—Why?—*Oh yes, it's the same.*—And if I pour the two glasses (B₁ and B₂) into these three (C₁ + C₂ + C₃), is it the same?—*There's more in the 3.*—And if I pour it back into the 2?—*Then there'll be the same (B₁ + B₂) as there (A₂).*'

Here is an example of the second type:

Fried (6;5) agreed that A₁ = A₂. A₁ was poured into B₁ + B₂. 'Is there as much lemonade as orangeade?—*Yes.*—Why?—*Because those (B₁ + B₂) are smaller than that (A₂).*—And if we pour the orangeade (A₂) as well into two glasses (doing so into B₃ + B₄, but putting more in B₃ than in B₄), is it the same?—*There's more orangeade than lemonade.*'—(B₃ + B₄ thus seemed to him more than B₁ + B₂).

A minute later he was given A₁ half full, and A₂ only a third full. 'Are they the same?—*No, three's more here (A₁).*—(A₁ was then poured into several glasses C.) *It's the same now as there (A₂).*' He finally decided, however: 'No, it doesn't change, because it's the same drink (i.e. A₁ = C₁ + C₂ + C₃ + C₄ and A₁ < A₂).'

These two types of intermediary reactions are important and enable us to dismiss an objection that doubtless occurred to the reader in §2. Instead of concluding that the notion of conservation has its origin in quantification properly so called (itself the result of progressive co-ordination of the relations involved), could we not explain the absence of the notion as being due merely to failure to understand the question as referring to the quantity as a whole? The child might simply be comparing one level with another or one width with another, without considering the total quantity of liquid, but that would not necessarily prove that he was incapable of so doing. If this were so, as soon as the idea of the whole quantity made its appearance, the child would suddenly discover conservation; he would at once understand that the liquid remains the same since nothing is added to or subtracted from it. And indeed, when Edi and other children state, when first questioned, that (A₂) and (B₁ + B₂) are 'the same', they give the

impression that the difference between them and the children in §2 is due merely to the fact that they interpret the question differently. The correct solution would then be the result of a kind of immediate identification and there would be no need for a complex process of quantification. But the intermediary reactions of this second stage make it clear that this too simple interpretation is not valid. If the child hesitates, if he gives a correct answer when the variations are slight but does not assume conservation when the variation in shape is greater, it is obvious that he understands the question but is not convinced *a priori* of the constancy of the whole quantity.

This being so, how are we to interpret the progress shown by children at the second stage? The two conditions laid down in §2 as defining the transition from 'gross quantity' to true quantification are beginning to be fulfilled.

At this stage the child is attempting to co-ordinate the perceptual relations involved and thus to transform them into true, operational relations. Whereas the child at the first stage is satisfied that two quantities of liquid are equal if the two levels are the same, irrespective of the width of the containers, the child at the second stage tries to take the two relations into account simultaneously, but without success, hesitating continually between this attempt at co-ordination and the influence of the perceptual illusions. This reaction is already apparent in the most advanced children of the first stage, but generally speaking it is typical of the second period. Here we have some examples, of which the first belongs to the earlier stage:

Lac (5;6): 'Your brother Lucien has this orangeade (A, $\frac{1}{2}$ full). Pour the same amount for yourself into this glass (L).—(He filled L to a higher level than that in A.) *No, I've got too much* (he poured some back so that L was $\frac{1}{2}$ full, i.e. the same level as in A).—Are they the same?—*No* (bringing L nearer to A and saying to himself): *Who has the most?*—*Yes, who has the most?*—(He pointed to A): *It's that one, because it's bigger.*—But you must have as much as Lucien.—(He added a little to L and compared the two levels.) *It's too much.* (He poured back the contents of L and began again. He gave himself the same level as in A, then added a little more so that L was about $\frac{2}{3}$ full.) *Oh! it's too much! It's not the same.* (In order to arrive at an equal quantity in L and A he then made the levels the same.)—You think you have the same amount to drink like that?—*Yes.*—(A was then poured into L'.)—*Oh! it's more!* (greatly astonished.) Lac thus showed that he was still at the first stage, although his first reactions suggested the second stage.

Edi (6;4). Glass A was $\frac{1}{2}$ filled. 'Pour as much orangeade into this one (L) as there is there (A).—(He filled L to the same level as that in A.)—Is there the same amount to drink?—*Yes.*—Exactly the same?—*No.*—Why not?—*That one (A) is bigger.*—What must you do to have the

same amount?—*Put some more in (filling L).*—Is that right?—*No.*—Who has more?—*Me (pouring some back).*—*No, the other one has more (A).*—(He continued to add more and then pour some back, without reaching a satisfactory conclusion.)

Wir (7;0): 'Can you pour as much into this one (L) as there is in that (A)?—*As much?*—(He filled it to the same level.)—Are they the same?—*No.* (He added some to L till it was $\frac{1}{2}$ full and then compared the levels.) *No, it's too much.* (He made the levels equal again.)—Who will have most to drink?—*Mummy (A), because the glass is bigger (adding some to L).*—Now are they the same?—*No, I've got more (pouring some back).*—Are they the same now, or has one of you got more?—*Mummy (A), because she has a bigger glass.* (He added some to L.) *No, now I've got more.* (He poured some back and again equalized the levels.) *No, mummy has more now (no satisfactory solution).'*

In each of these cases, the child begins, as at the first stage, by filling the narrow glass L to the same level as the wide glass A. But unlike the earlier children, he then discovers, by comparing the two columns of equal height, that one is wider than the other and decides that the first glass contains more because it is bigger. Thus a second relation, that of the width, is explicitly brought into the picture and 'logically multiplied' (see Glossary) with that of the levels. In order to arrive at equality, the child pours a little more liquid into glass L, thus proving the reality of this multiplication of relations. But as soon as the level of the liquid in the narrow glass is higher than that in the wide one, the child forgets the cross sections and thinks that L contains more than A, thus showing clearly the difficulties of this multiplicative operation. When he is concerned with the unequal levels, he forgets the width, and when he notices the difference in width he forgets what he has just said about the relation between the levels. Hence it is only when the levels are equal that he attempts a logical multiplication of the relations of height and width, and when the attempt has been made, one or other of the relations becomes all-important and he is left hesitating between them.

It is obvious, however, that even if the operation of logical multiplication of relations were carried through by the child of this stage it would not suffice for the construction of conservation of the whole quantity unless the height and width were simply permuted. A column of liquid whose height increases and whose width diminishes with respect to another column may be greater, equal, or less in volume than the other. In order to be certain that there is equality, the intensive graduation must be completed by an extensive quantification, i.e., it must be possible to establish a true proportion, and not merely a qualitative correlation, between the gain in height and the loss in width. In other words, there must be partition of some kind to supplement the co-ordination.

Now it is during the second stage, and in close connection with the logical co-ordination just described, that the child begins to understand that a whole remains identical with itself when it is divided into two halves. Edi and Pie, examples of the first stage, stated this fact when A₁ was poured into B₁ + B₂, but this understanding of partition is short-lived and discontinuous, like the multiplication of relations. As soon as B₁ and B₂ are poured into C₁ + C₂ + C₃, Edi and Pie no longer think there is conservation. 'In three there's more', they say, and Edi even goes so far as to think that the more a given quantity is sub-divided the more there will be of it.

In conclusion, then, multiplication of relations and partition go hand in hand. Both of them make their appearance and begin to develop during this second stage, both are halted as a result of the same limitations. What then is the link that unites these two types of operation? The analysis of the third stage will provide the answer.

§4. Stage III: Necessary conservation

In the replies characteristic of the third stage children state immediately, or almost immediately, that the quantities of liquid are conserved, and this irrespective of the number and nature of the changes made. When the child discovers this invariance, he states it as something so simple and natural that it seems to be independent of any multiplication of relations and partition. Our problem is therefore to discover whether this independence is real or only apparent, and if the latter is the case, to determine how the various intervening factors are linked.

Here are some examples:

Aes (6;6). A₁ and A₂ were $\frac{3}{4}$ filled, and then A₁ was poured into P₁, which was wide and low. 'Is there still as much orangeade as there was in the other glass?—*There's less.*—(A₂, which was supposed to be his glass, was poured into P₂.) Will you still have the same amount to drink now?—*Oh yes! It's the same, It seems as if there's less, because it's bigger (=wider), but it's the same.*—(P₁ and P₂ were poured back into A₁ and A₂, and A₁ was then poured into B₁ + B₂.) Has Roger got more than you now?—*He's got the same (definitely).*—And if I pour yours into 4 glasses (A₂ into C₁ + C₂ + C₃ + C₄)?—*It'll still be the same.'*

Geo (6;6). Her glass was A₁, $\frac{1}{2}$ full, and A₂, only $\frac{1}{3}$ full was supposed to be Madeleine's. 'Who has more?—*I've got more.*—That's right. But Madeleine wants to have the same amount. She divides hers by pouring it into two glasses (C₁ + C₂) and says: "Now I've got more, or at any rate the same amount as you."—Who has more now?—(After some thought) *It's still me.*—She then pours it into 3 glasses (C₁ + C₂ + C₃). Who has more now?—*Still me.*—Then she pours it into a lot of glasses (C₁, C₂ and C₃ were poured back into A₂, the contents of which

were then divided between 6 little glasses C). Who has more now?—*Madeleine has more, because it's been poured into the other glasses.*—And if we put it all back into here (A_2) where will it come up to?—(She reflected.) *No, Madeleine has less. I thought she had more, but she hasn't.*—Can't it be more?—*No.*—(Glasses C were poured back into A_2 , then A_2 was poured into 8 little glasses.) And now?—*No, it's still the same. It's the same all the time.* Finally she was given two new glasses A_3 and A_4 , both half-full, and A_3 was poured into $B_1 + B_2$: '*She has the same.*—Are you sure it's the same?—*Yes, it's only been poured out.*'

Bert (7;2): 'The orangeade (A_1 , $\frac{3}{4}$ full) is for Jacqueline, the lemonade (A_2 , $\frac{1}{2}$ full) is for you. Who has more?—*Jacqueline.*—You pour yours (A_2) into these two ($B_1 + B_2$, which were then full). Who has more?—*It's still Jacqueline.*—Why?—*Because she has more.*—And if you pour this (B_1) into those ($C_1 + C_2$)?—*It's still Jacqueline, because she has a lot.*—Every change produced the same result: '*It's Jacqueline, because I saw before that she had more.*' Then A_3 , equal to A_4 , was poured into $C_1 + C_2$: '*It's still the same, because I saw before in the other glass that it was the same.*—But how can it still be the same?—*You empty it and put it back in the others!*'

Eus (7;2). A_1 was $\frac{3}{4}$ full and A_2 was $\frac{1}{2}$ full. A_2 was poured into $C_1 + C_2$: 'Are they the same now?—*No, You've poured it out of the same glass (A_2). Like that you can never make them the same.*' Then with A_1 equal to A_2 , A_2 was poured into $B_1 + B_2$, etc.: '*It's still the same, because it still comes from the same glass.*'

These few examples are sufficient to show clearly which of the hypotheses in §3 corresponds to what actually occurs. If we merely consider the answers of Bert and Eus, who are 7, it seems that the global comparison between the initial and final state of the liquid is sufficient to enable the child to see that there is conservation, independently of any multiplication of relations or partition. 'It's still the same', Eus says, 'because it always comes from the same glass'. At a certain level of development, therefore, the notion of conservation seems to be the result of an *a priori* analytic deduction (see Glossary) which makes it possible to dispense with observation of the relations and even with the experiment itself. If, however, we examine the replies of Aes and Geo, who still hesitate for a moment before being quite certain, the mechanism of their construction becomes apparent, and we are compelled to recognize that the reasoning that leads to the affirmation of conservation essentially consists in co-ordination of the relations, with its two-fold aspect of logical multiplication of relations and mathematical composition of parts and proportions. Aes, for instance, begins by thinking that the liquid in glass A becomes less when poured into glass P, but immediately afterwards says: 'It seems as if there's less, because it's bigger (=wider), but it's the same.' In other words, he corrects his mistake by co-ordinating the relations of

height and width. In general, when children of this stage are confronted with the problem of glasses A and L, their answers show, unlike those of the earlier stages, correct co-ordination of the relations involved.

Aes (6;6). Although he began by pouring into L (long and narrow) a column of liquid the same height as in A in order to obtain the same quantity, he very soon corrected himself: 'Is it the same?—*Yes, it's the same height. . . . Oh no! that one (L) is narrower and this one (A) is wider (and he added more to L).*'

Geo (6;6) at once filled glass L $\frac{3}{4}$ full to equal $\frac{1}{2}$ in glass A: 'Is that right?—*It's about right.*—Is there the same amount to drink?—*Yes, it's the same.*—Why?—*Because it's narrower here (L) and wider there (A).*—What can you do to be sure it's the same (giving her the glasses)?—(She took glass A_2 and poured into it the liquid in L: it was approximately the same quantity as in A_1 .)'

Bert (7;2) began by filling L to the same level as in A, then added some more 'because the glass is smaller: you think it's the same, but it's not true.'

Eus (7;2) at once poured into L a column of liquid higher ($\frac{3}{4}$) than in A ($\frac{1}{2}$) and explained his action by saying: '*Here (A), it's lower, but it's the same as there (L).*'

Ela (7;0): '*In this one (L) we must put more, because it's narrower, and in the other there's more room because it's wider.*'

These children, all of whom show in other experiments that they recognize conservation of quantity, succeed without difficulty in the present experiment in multiplying the relations of height and width in glasses A and L. It should be stressed that this question of the relation between A and L was put to all our children before the questions relating to conservation of quantities. It is not, therefore, the discovery of conservation which makes possible the multiplication of relations, but the contrary. Moreover, this question appears to be on the whole rather easier than that of conservation in general, since correct answers are given slightly earlier than those assuming invariance. This fact constitutes a further reason for assuming that conservation of quantities, even when it is affirmed categorically, as though it was an *a priori* judgement, presupposes a much more complex construction than would at first appear.

It is obvious that logical multiplication of relations is not sufficient to ensure the discovery of the invariance of whole quantities, and we shall now see why. When the child, after estimating the quantities from one point of view only, begins to co-ordinate the relations one with another, he does indeed construct a multi-dimensional whole, but it is a whole that remains 'intensive' and is not susceptible of 'extensive' measurement so long as the logical

multiplication is not supplemented by truly mathematical considerations.

What, in fact, is logical multiplication of the relationships of height and width? If we have a series of containers of the type A (A_1, A_2, A_3 , etc.) containing liquid at increasing levels, we shall say that the child can add relations if from $A_1 < A_2$, and $A_2 < A_3$, he can deduce that $A_1 < A_3$, the operation obviously taking place only on the practical plane, the levels being clearly perceptible. If we take another series of containers of increasing width (L, B, A, P, etc.) we shall again say that there is addition if there is uni-dimensional co-ordination of the relationships. But we shall say that there is logical multiplication of relationships when the child compares the containers from the point of view of the two relationships *simultaneously*, e.g., when the column of liquid in L is both higher and narrower than that in A.

These logical multiplications of relationships are essential to the child's solutions, since he cannot assign any numerical value to the two dimensions and is therefore unable to multiply them arithmetically. Moreover, this logical operation makes possible the conception of a further relationship, that of total quantity, which is the logical product of height and width. For instance, if A is one-fifth full and P is quite full, the liquid in P being at the same time higher and wider than that in A, no child will have any hesitation in saying that P contains more than A. When Geo compares A_1 , which is half full with A_2 , one-third full, he at once deduces that since the cross-sections are equal there is more liquid in the container in which the level is higher. In a word, logical multiplication of relationships is the necessary intermediary between gross, uni-dimensional quantity and extensive quantification, which we shall consider presently.

But obviously these elementary operations can only lead to simple seriations or 'intensive' graduations, since the only possible conclusions are the following: (1) if the two relationships vary in the same direction, or (2) if one remains constant and the other varies, or (3) if both remain constant, then it is possible to discover whether the total quantity increases, decreases, or remains the same. If, on the other hand, the height increases and the width decreases, or conversely, it is impossible to know whether the total quantity increases, decreases, or remains constant. In general, the child will therefore be able to grasp increasing or decreasing series in certain conditions, but he will not be able to tell by how much one quantity is greater or less than another, or whether the total quantity increases or decreases when the relationships that intervene vary in opposite directions.

Now the notion of conservation of wholes, which is acquired by

children at this stage, presupposes a quantification wide enough to cover the case in which elementary relationships vary in opposite directions, and it therefore presupposes the discovery of 'extensive' quantities. Indeed, if the child is to assume conservation, he must not only be aware that when the width and height of the columns are the same the total quantity remains constant, but also that it remains constant in spite of the fact that the height increases and the width decreases. But this conclusion is impossible within the limits of logical multiplication of relationships. How then does the child succeed in going beyond these limits if he has no numerical data or measurements in the true sense? This is the problem that we must consider, and it is bound up with the whole question of the transition from intensive to extensive quantity.

We might, of course, at this point, confine ourselves to saying, like Geo, that there is conservation because 'it's only been poured out', or like Eus, because 'it still comes from the same glass'. Conservation would then be identified with mere logical permanence, without the intervention of mathematics. But in our view, it then becomes legitimate to ask why the child does not make this discovery before he reaches the third stage. Children of 4 or 5 know as well as older children that 'it's only been poured out' or that 'it still comes from the same glass', and yet for them the quantity varies. How is it that they do not see the final and initial states as being identical, whereas at 6 or 7 they will find no difficulty in doing so?

The reason is that here a second process intervenes, one which is synchronic but distinct from the earlier one, its relations with which need careful examination, since they dominate the whole development of mathematical notions. It is the intervention of the notion of the 'unit', i.e., extensive quantification in the form either of arithmetical partition, or, which comes to the same thing, of proportions.

To take a concrete example, when Aes tried to put into L a quantity of liquid equal to that in A (one-fifth full), he poured in a higher column and said 'it's the same . . . because this one (L) is narrower and that one (A) is wider'. What does this mean? If he was merely using logical multiplication, he could only conclude that the liquids in A and L were equal if the height of the one and the width of the other were interchangeable. Something more is therefore implied in his reasoning, the feeling of a definite proportion such that L lost in width what it gained in height, or that the height of A was to its width as the height of L was to its width, or, to put it more simply, that the increase in height when A was poured into L was equivalent to the decrease in width.

At the first stage, the child confines himself to establishing simple, qualitative differences. During the following stages, when he is using only logical multiplication of relationships, he progressively grades these differences, in one or more dimensions, into 'intensive' seriations of two or more terms. But these series, except when there is complete equality, involve only asymmetrical relationships of difference. Since these differences can be seriated, they result in intensive quantification, but when two given quantities show two relationships of difference, there is no means of equating them. In other words, multiplication of relationships is a multi-dimensional seriation leading only to further seriations, and in no way allowing of the division of a given quantity into units that are recognized as equal and yet distinct. On the other hand, both proportion and numerical partition imply the fusion of asymmetrical relationships of difference with those of equality, and it is this combination of equalities and differences, or, to put it more briefly, this equating of differences, which constitutes the transition from intensive to extensive quantity, and explains the arithmetization of logical multiplication.

We shall now attempt to express what occurs in terms of actual operations that are psychologically real. First of all, it is clear that the child would have no means of gauging the equality or non-equality of the various quantities in A_1 and $(B_1 + B_2)$ or in P, L , etc., if he were merely asked to compare them. The fact that the liquid is poured from one container into another does, of course, suggest equality, but as we have seen, this action is not sufficient to explain conservation, since the younger children think that the change in shape involves a change in quantity. The action of pouring does, however, lead to the notion of invariance of the quantity when it is supported by the following operations. The empty glass A is filled to one-fifth. When A is poured into L , the liquid in L will be higher but narrower. While the child is still on the plane of qualitative seriation, he is capable of co-ordinating two relationships of level, or width, or both together. This means that, on comparing the liquids in A and L he can at once see that the level in L is higher than that in A , i.e. that it is equal to that in A plus a difference. Similarly, he can compare the liquids and see that L is narrower than A , the width of L thus being equal to that of A , less a difference. But there is no means, in these simple comparisons, of quantifying these relationships otherwise than as 'more' or 'less'. Our contention is that at a given moment the child grasps that the differences compensate one another, and this is the beginning of extensive quantification, because then two heterogeneous qualitative relationships (increase in level and decrease in width) are seen to be equal, though still preserving

their value of asymmetrical difference. It is thus that proportion comes into being, through the combination of equality and asymmetrical relationship.

In a sense, this proportion is already a partition, for the equating of the increase in level with the decrease in width means that the quantity as a whole is no longer viewed as a qualitative totality whose value changes with each change of shape, but is structured as a sum susceptible of division into units. The criterion is the following. There is arithmetical partition as soon as the elements of a whole can be equated with one another and yet remain distinct.

Moreover, the equating of differences, which we have just stated to be the principle of extensive quantification, gives rise during the third stage to truly numerical partition, which is not only synchronic but complementary to the discovery of proportions. Thus for Aes, it is obvious that when A_1 is poured into $2B$ or $4C$ it still remains A_1 , and although Geo is at first reluctant to accept $6C$ as being equal to A , when he finally does so he generalizes and assumes that the same is true of $8C$; etc. During the first stage, and during the second except in very simple cases of partition, this was not so.

In order to explain this evolution we must understand that, for instance, half a quantity is not only a quantity which is equal to another quantity and which when added to it constitutes the original quantity, but also that it is equal to the difference between the whole and the other half. Without this second condition, the relationship between the half and the whole could not be understood and the notion of the whole would cease to exist after the division. Numerical partition is therefore essentially an equating of differences, like proportion itself, but in the case of $A = B_1 + B_2$, the two halves B_1 and B_2 are seen as equal, whereas in the case of $A = L$, it is only the differences that are equated.

In conclusion, we have shown the extreme simplicity, at bottom, of the process of quantification as revealed in the child's discovery of the conservation of quantities. He begins—and makes no further progress during the first stage—by considering only unco-ordinated perceptual relationships of qualitative equality or difference, thus acquiring the notions of gross quantities and qualities not susceptible of composition. During the second stage, there begins a process of logical co-ordination, which is completed at the third stage and leads to classification when there is equality and to seriation when there is difference, this seriation resulting in intensive quantities. Finally, the third stage is characterized by the construction of extensive quantities through the

equating of intensive differences and therefore through the arithmetization of logical groupings (see Glossary).¹

¹ For the sake of simplicity, we have here restricted ourselves to explaining the discovery of conservation of quantities of liquid in terms of the proportion established by the child between differences in height and width of columns of water, this being the method used by our subjects. It would obviously be possible also to envisage a purely logical (non-arithmetical) conservation in cases in which the displaced parts could be abstracted individually, or in which the differences in height and width could be compensated merely by substitution, height becoming width and vice versa.

CHAPTER II

CONSERVATION OF DISCONTINUOUS
QUANTITIES AND ITS RELATION TO
ONE-ONE CORRESPONDENCE¹

THE experiments described in the previous chapter can all be repeated with (discontinuous quantities) that can be evaluated globally when the elements are massed and counted when they are separated. (Sets of beads, for instance, can be used.) If they are put into the containers used in Chapter I, they can serve for the same evaluations as the liquids (level, cross-section, etc.), and in addition they are material for a further global quantification with which children are familiar: that of the length of necklaces made from the beads. The evaluation of this length can thus be used in each case to check the quantification of the contents of the various containers used. On the other hand, when the beads are considered as separate units they can be used in operations of correspondence. If the child is told, for instance, to put beads into a container, one by one, at the same time as the experimenter is putting beads one by one into another container, he can then be asked whether the total quantities are the same, with or without identity in the shape of the two containers.

In going on from the analysis of continuous quantities to that of discontinuous quantities we are therefore not merely checking our earlier findings. We are also making a preliminary study of the relationship between conservation of quantities and the development of one-one correspondence, which is, as is well known, one of the origins of number. We shall then be in a better position to approach the question of cardinal and ordinal correspondence as such.

It should be noted that the stages we shall find here correspond exactly to those of the previous chapter.

§1. Stage I: Absence of conservation

During the first stage there is no conservation of the sets of beads, just as there was no conservation of the quantity of liquid. The child not only thinks that the total quantity changes when a

¹ With the collaboration of M. Juan Jaen.